

**CATASTROPHIC
SPORTS INJURY RESEARCH**

THIRTY-SIXTH ANNUAL REPORT

FALL 1982 - SPRING 2018

**From the
National Center for Catastrophic Sport Injury Research
At The University of North Carolina at Chapel Hill**

Website: nccsir.unc.edu

Prepared by:
Kristen L. Kucera, MSPH, PhD, LAT, ATC
University of North Carolina
Chapel Hill, NC 27514

Robert C. Cantu, MD
Medical Director, National Center for Catastrophic Sport Injury Research
Emerson Hospital
Concord, MA 01742

FINAL
October 3, 2019

DO NOT DISTRIBUTE
Report #: 2019-03



Acknowledgements:

We acknowledge the significant contributions of recently retired Frederick O. Mueller, Ph.D. who directed The National Center for Catastrophic Sport Injury Research (NCCSIR) from 1982 to 2013. Dr. Mueller's work during those 30 years has improved the safety of football for the participants and these impacts are demonstrated in the pages of this report.

We also acknowledge NCCSIR staff members Courtney Haley and Lily Wang and members of the Consortium for Catastrophic Sport Injury Monitoring: Drs. Douglas Casa, Jonathan Drezner, Kevin Guskiewicz, Johna Register-Mihalik, Steve Marshall, Dawn Comstock, David Klossner, Tom Dompier, Zack Kerr, Erin Wasserman, and Christine Collins.

We also thank all the athletes, families, coaches, athletic trainers, medical providers, school staff, state associations, researchers, journalists, and others who have participated in this research and have shared information with the NCCSIR.

Funding & Disclosures:

The National Center for Catastrophic Sport Injury Research is supported by the American Football Coaches Association (AFCA), the National Collegiate Athletic Association (NCAA), the National Federation of State High School Associations (NFHS), the National Athletic Trainers' Association (NATA), the American Medical Society for Sports Medicine (AMSSM), the National Operating Committee on Standards for Athletic Equipment (NOCSAE), and The University of North Carolina at Chapel Hill (UNC-CH).

All rights reserved. This material may not be published, broadcast, rewritten or redistributed in whole or part without express written permission. Contact the National Center for Catastrophic Sport Injury Research for all questions regarding this report at nccsir@unc.edu.

TABLE OF CONTENTS

	<u>Page</u>
Introduction	1
Methods	
Outcome Definitions	2
Data Collection	2
Participation Data	3
Analysis	4
Results	
Current year summary – AY2017-2018	4
Overall summary	5
Discussion	7
Recommendations	8
Case Summaries for AY2017-2018	12
References	22

LIST OF TABLES

	<u>Page</u>
Table 1: Number of All catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)	23
Table 2: Number of Direct catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)	24
Table 3: Number of Indirect catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)	25
Table 4a: Number of Direct catastrophic injuries/illnesses by severity by sport: High school all years combined	26
Table 4b: Number of Indirect catastrophic injuries/illnesses by severity by sport: High school all years combined	27
Table 5a: Number of Direct catastrophic injuries/illnesses by severity by sport: College all years combined	28
Table 5b: Number of Indirect catastrophic injuries/illnesses by severity by sport: College all years combined	29
Table 6a: Number of Catastrophic injuries/illnesses by Severity by year: High school	30
Table 6b: Number of Catastrophic injuries/illnesses by Severity by year: College	31
Table 7a: Rate of direct catastrophic injuries/illnesses by severity by year: High school	32
Table 7b: Rate of direct catastrophic injuries/illnesses by severity by year: College	33
Table 8a: Rate of indirect catastrophic injuries/illnesses by severity by year: High School	34
Table 8b: Rate of indirect catastrophic injuries/illnesses by severity by year: College	35
Table 9a: Rate of Direct catastrophic injuries/illnesses by level and severity by sport: High school	36
Table 9b: Rate of Direct catastrophic injuries/illnesses by level and severity by sport: College	37

Table 10a: Rate of Indirect catastrophic injuries/illnesses by level and severity by sport: High school	38
Table 10b: Rate of Indirect catastrophic injuries/illnesses by level and severity by sport: College	39
Table 11: Characteristics of all catastrophic injuries/illnesses AY 2017/18	40
Table 12: Participation numbers, 1982-1983 to 2017-2018	44

Note: Tables I-IX are located in a separate Appendix document located here: <http://nccsir.unc.edu/reports/>.

LIST OF FIGURES

	<u>Page</u>
Figure 1: Rates of <i>fatal</i> catastrophic direct and indirect injuries/illnesses by sport-gender among <u>high school</u> participants, 1982/83-2017/18	45
Figure 2: Rates of <i>all</i> catastrophic direct and indirect injuries/illnesses by sport-gender among <u>high school</u> participants, 1982/83-2017/18	46
Figure 3: Rates of <i>fatal</i> catastrophic direct and indirect injuries/illnesses by sport-gender among <u>Collegiate</u> participants, 1982/83-2017/18	47
Figure 4: Rates of <i>all</i> catastrophic direct and indirect injuries/illnesses by sport-gender among <u>Collegiate</u> participants, 1982/83-2017/18	48

INTRODUCTION

In 1931, the American Football Coaches Association (AFCA) initiated the First Annual Survey of Football Fatalities and this research has been conducted at the University of North Carolina at Chapel Hill since 1965. In 1977, the National Collegiate Athletic Association (NCAA) initiated a National Survey of Catastrophic Football Injuries, which is also conducted at the University of North Carolina. As a result of these research projects important contributions to the sport of football have been made. Most notable have been the 1976 rule changes making it illegal to make initial contact with the head and face while blocking and tackling, the National Operating Committee on Standards for Athletic Equipment (NOCSAE) football helmet standard, improved medical care for the participants, and better coaching techniques.

Due to the success of these two football projects the research was expanded to all sports for both men and women, and a National Center for Catastrophic Sports Injury Research (NCCSIR) was established in 1982. The decision to expand this research was based on the following factors:

1. Research based on reliable data is essential if progress is to be made in sports safety.
2. The paucity of information on injuries in all sports.
3. The rapid expansion and lack of injury information in women's sports.

In 1987, a joint endeavor was initiated with the Section on Sports Medicine of the American Association of Neurological Surgeons. The purpose of this collaboration was to enhance the collection of medical data. Dr. Robert C. Cantu, Chairman, Department of Surgery and Chief, Neurosurgery Service, Emerson Hospital, in Concord, MA, is the Medical Director of the NCCSIR and has been responsible for evaluating the medical data. Dr. Cantu is also a Past-President of the American College of Sports Medicine. The NCCSIR was directed for 30 years by Dr. Frederick Mueller. Dr. Mueller retired in the Spring of 2013 and the NCCSIR continues under new direction (Dr. Kucera). The NCCSIR has expanded to become a consortium of universities (University of North Carolina, Boston University, University of Washington, University of Connecticut, University of Colorado, University of Maryland) with expertise in head/neck, cardiac, and heat-related sports medicine (these three areas account for the overwhelming majority of catastrophic events).

To learn more about NCCSIR please visit: <http://nccsir.unc.edu/about/>

To learn more about the Consortium please visit: <http://nccsir.unc.edu/consortia-and-partners/>

To access online reports please visit: <http://nccsir.unc.edu/reports/>

METHODS

Outcome Definitions

For the purpose of this research the term catastrophic is defined as any severe injury incurred during participation in a school/college sponsored sport. Catastrophic is divided into the following three definitions:

1. **Fatality**
2. **Non-Fatal** - permanent severe functional disability.
3. **Serious** - no permanent functional disability but severe injury. An example would be fractured cervical vertebra with no paralysis.

Sports injuries are also considered traumatic (or direct) or exertional/systemic (or indirect). The definition for direct and indirect is as follows:

Direct - Those injuries that resulted directly from participation in the skills of the sport.

Indirect - Those injuries that were caused by systemic failure as a result of exertion while participating in a sport activity or by a complication that was secondary to a non-fatal injury.

Note: Beginning in 2014, NCCSIR also collects non sport-related injuries such as sudden cardiac arrest that occurred outside of sport activity (e.g., during sleep). These events were not included in the tables but are described in the Case Summary sections.

Data Collection

Data were compiled with the assistance of coaches, athletic trainers, athletic directors, executive officers of state and national athletic organizations, online news reports, and professional associates of the researchers. Data collection would not have been possible without the support of the NCAA, the National Federation of State High School Associations (NFHS),

and the AFCA. Upon receiving information concerning a possible catastrophic sports injury, contact by telephone, email or personal letter and questionnaire was initiated with the injured player's athletic trainer, athletic director, or coach. Data collected included background information on the athlete (age, height, weight, experience, previous injury, etc.), accident information, immediate and post-accident medical care, type injury, and equipment involved. Autopsy reports are used when available. In order to improve overall capture of catastrophic sport injury and illness events, NCCSIR and the Consortium for Catastrophic Injury Monitoring in Sport developed an online portal where anyone can report a catastrophic event: <https://www.sportinjuryreport.org>. The portal was activated in January 2015.

Participation in Sport

Yearly participation estimates for high school athletes are obtained from NFHS participation reports (available online: <https://www.nfhs.org/ParticipationStatistics/ParticipationStatistics/>). NFHS high school annual athletic participation for 2017/18 included 7,677,337 athletes (4,421,063 males and 3,256,274 females). Yearly participation estimates for collegiate level athletes are obtained from the National Collegiate Athletic Association (NCAA) participation reports (accessed online: https://ncaaorg.s3.amazonaws.com/research/sportpart/Oct2018RES_2017-18SportsSponsorshipParticipationRatesReport.pdf). NCAA participation for 2017/18 in championship sports was 494,992 athletes. There were 278,614 males and 216,378 females. There were also 3,314 males in non-championship sports (archery, badminton, bowling, equestrian, rowing, rugby, sailing, and squash) and 2,427 females participating in emerging sports (archery, badminton, equestrian, rugby, squash, synchronized swimming, team handball, and triathlon).

During the entire 35 year period from the fall of 1982 through the spring of 2018, there were 228,610,552 high school participant-seasons in the sports covered by this report and approximately 13,069,182 college participant-seasons (Table 12).

Not all high schools and colleges are members of the NFHS and NCAA. Complete data is not available for the non-member schools. Therefore, these participation numbers underestimate the total number of high school and collegiate participants in the United States.

Analysis

Frequencies and incidence rates of catastrophic injury per 100,000 participants were calculated over the entire 36-year period and stratified by level (high school and college) and sport. Incidence rates were stratified by direct versus indirect and by severity. **Note: if there were no events in the sport for a particular year, the year is excluded from the frequency Table. Rates with number of incidents less than 5 should be interpreted with caution.**

It is important to note that information is continually being updated due to the fact that catastrophic injury information may not always reach the NCCSIR in time to be included in the current final report. The report includes data that is reported to the NCCSIR by the NCAA, the NFHS, online reports, colleagues, coaches, and athletic trainers. There may be additional catastrophic injuries that are not reported to the NCCSIR. The authors acknowledge that not every catastrophic injury is included in this report.

RESULTS

Current AY2017-2018 Summary

From July 1, 2017 to June 30, 2018 there were a total of 99 catastrophic injuries/illnesses captured by NCCSIR among high school and college organized sport participants. Of these, 85 events were due to or occurred during sport-related activities (Table 11). There were also 14 catastrophic events that occurred during non-sport related activity. All were cardiac-related.

Sport-related events: The majority of the sport-related catastrophic events (n=85) were at the high school level (78%, n=66). Member institutions for collegiate cases included NCAA and National Association of Intercollegiate Athletics (NAIA). Overall 24.7% of cases were fatal, 9.4% were nonfatal, 60.0% were serious with recovery, and 5.9% were unknown. Forty-six percent (n=39) were due to direct (traumatic injury) causes and over half occurred in competition (50.6%) followed by practice (29.4%). The majority of events occurred to athletes participating in the following sports: football (54.1%), basketball (15.3%), soccer (9.4%), baseball (4.7%), cross country (3.5%), lacrosse (3.5%), track and field (2.4%), wrestling (2.4%), swimming (2.4%), cheerleading (1.2%) and ice hockey (1.2%). Areas of the body most commonly affected

were heart (38.8%), spine (21.2%), and head/brain (15.3%). Sudden cardiac arrest (38.8%) was the most common types of events followed by brain trauma (14.1%) and other traumatic injuries (9.4%).

Direct events: 12.8% of direct events were fatal, 20.5% non-fatal, 53.8% serious with recovery, and 12.8% unknown. A greater proportion of direct events occurred in competition versus practice (69.2% versus 25.6%) and were due to contact with another player (56.4%) or apparatus/object (12.8%). The highest proportion was to the spine (46.2%) and head/brain (33.3%) followed by other traumatic injury (15.4%) and commotio cordis (5.1%). The majority occurred in football (79.5%) followed by soccer (5.1%) and baseball (5.1%).

Indirect events: 34.8% of indirect events were fatal and 65.2% were serious with recovery. Similar proportions of indirect events occurred in competition and practice (34.8% versus 32.6%) and 10.8% occurred during conditioning and weight training sessions. The majority were cardiac-related (71.7%), followed by heat related (15.2%) and rhabdomyolysis and other (2.2% each). Football (32.6%) and basketball (28.3%) comprised the majority, followed by soccer (13.0%) and track and field, baseball, lacrosse, cross country, and swimming (4.3% each).

Overall Summary

During this 36-year period, there were 2,686 catastrophic sport-related injuries/illnesses at high school and college levels (Table 1 – excluding cheerleading, drill team, and rodeo there were 2,566). The majority were non-fatal (64%) and from traumatic or direct mechanisms (65%), and among high school participants (79%). The proportion of fatal (38% versus 35%) and direct (65% versus 62%) were not different by high school compared to college level.

The 85 sport-related catastrophic injuries and conditions captured in 2017/18 is not different from the previous year (80 in 2016/17) and a 26% decrease from 2015/16—a result of a lower number of traumatic brain and cervical spine/spinal cord injuries in 2017/18 and 2016/17. *Note: see limitations regarding the interpretation of this percentage difference.*

Traumatic Injuries (Direct) by Sport: For high school sports, football had the highest *number* of traumatic injuries (direct) catastrophic events, followed by female cheerleading, wrestling, baseball, and male track and field (Table 4a). Accounting for the number of participants in the sport, male and female cheerleading, male gymnastics, football, and male ice hockey had the highest rates per 100,000 participants (Figure 2, Table 9a). Similar results were observed when restricted to fatal events (Figure 1).

For college sports, football had the highest *number* of traumatic injuries (direct) catastrophic events, followed by female cheerleading, baseball, and male track and field (Table 5a). Accounting for the number of participants in the sport, male gymnastics, female skiing, football, male ice hockey, male skiing, equestrian and female gymnastics had the highest rates per 100,000 participants (Figure 4, Table 9b). Similar results were observed when restricted to fatal events (Figure 3).

Exertional and Systemic Conditions (Indirect) by Sport: For high school sports, football had the highest *number* of exertional and systemic (indirect) catastrophic events, followed by male basketball, male track and field, male soccer, wrestling, baseball, and male cross country (Table 4b). Accounting for the number of participants in the sport, rowing, male basketball, football, male ice hockey, male water polo, and male lacrosse had the highest rates per 100,000 participants (Figure 2, Table 10a). Similar results were observed when restricted to fatal events (Figure 1).

For college sports, football had the highest *number* of exertional and systemic (indirect) catastrophic events, followed by male basketball, wrestling, female basketball, baseball, male soccer, and male swimming (Table 5b). Accounting for the number of participants in the sport, male basketball, male water polo, male skiing, football, male wrestling, male ice hockey, male swimming, and male rowing had the highest rates per 100,000 participants (Figure 4, Table 10b). Similar results were observed when restricted to fatal events (Figure 3).

DISCUSSION

The following strengths and limitations should be noted:

- Data have been collected by The National Center for Catastrophic Sport Injury Research for all high school and college sports since 1982 using consistent definitions and methodology over a 30+ year period. These data are provided annually to sport organizations (NCAA, NFHS, AFCA), researchers and the public. Sports medicine advisory committees, sport rules committees, and coaching committees review the reports and have used these data to inform and evaluate safety recommendations, medical care, and rule changes.
- Catastrophic events are primarily captured through publicly available media reports. Therefore, not all catastrophic events are captured. Particularly, for non-fatal catastrophic events, which may not be reported in the media as comprehensively as fatalities. Under-reporting may also be due to outcome definitions used (e.g. timing of the event) and event locations (e.g. at home, personal conditioning). **In order to improve overall capture of these events, NCCSIR and the Consortium for Catastrophic Injury Monitoring in Sport have developed an online portal where anyone can report a catastrophic event: <https://www.sportinjuryreport.org>.** The online portal was activated in January 2015. Any observed changes in annual number of events may be attributed to these described improvements in data collection methods.
- Details surrounding catastrophic events that are only captured through publicly available media reports may not be completely accurate in the absence of the actual autopsy or medical reports.
- Incidence rates were calculated using participation estimates from NFHS and the NCAA in the rate denominator (Table 12). These participation estimates do not include schools that are not members of these two associations. Participation data were not available for these non-member schools. At present NFHS and NCAA are the only estimates available. Therefore, the participation numbers (rate denominator) in this report are underestimated, which results in an overestimate of the actual incidence rate.
- It is important to note that catastrophic events are rare and statistical power for some strata comparisons are limited. Rates with number of incidents less than 5 should be interpreted with caution.

RECOMMENDATIONS

1. Each athlete should have a complete physical examination with a medical history and an annual health history update.
2. All personnel involved with training athletes should emphasize proper, gradual, and sport-specific physical conditioning.
3. Every school should strive to have a certified athletic trainer.
4. Each school should have a written emergency action plan (EAP) in place, all personnel should have copies, and procedures should be reviewed and practiced annually.
 - The Centers for Disease Control and Prevention (CDC) has guidelines and templates for these plans (<http://www.cdc.gov/niosh/docs/2004-101/emrgact/emrgact1.html>).
 - NCAA and the NFHS have guidelines for these plans at the following websites: www.nfhs.org and www.ncaa.org.
 - **An automated external defibrillator (AED) should be available and accessible onsite and medical and coaching staff should be trained in the use.**
5. There should be an emphasis on employing well trained athletic personnel, providing excellent facilities, and securing the safest and best equipment available.
6. There should be strict enforcement of game rules and administrative regulations to protect the health of the athlete and reduce the risk of catastrophic injury. Coaches and school officials must support the game officials in their rulings during the sporting event.
7. Coaches should be educated on and have the ability to teach the proper fundamental skills of the specific sport. Specific to football, the proper fundamentals of blocking and tackling should be emphasized to help reduce head and neck injuries, especially with keeping the head out of blocking and tackling.
8. Weight loss in wrestling to make weight for a match can be dangerous and cause serious injury or death. Coaches should be aware of safety precautions and rules associated with this practice.
9. There should be continued surveillance and safety research in athletics (rules, facilities, equipment, medical care and procedures).
10. **Sudden cardiac arrest:** The number of exertional and systemic (indirect) cardiac related events has increased over the years and it is recommended that schools have and emergency action plan and automated external defibrillators (AED) available and

accessible on-site for emergency situations. Early detection and defibrillation is critical for survival (3-5 minutes recommended). (Casa et al. 2012)

- See also Drezner et al. 2007 for additional information about sudden cardiac arrest preparedness and management: <http://www.nata.org/sites/default/files/sudden-cardiac-arrest-consensus-statement.pdf>

11. **Heat-illness:** All personnel associated with sport participation should be cognizant of the safety measures related to physical activity in hot weather. Heat stroke and heat exhaustion are prevented by careful control of various factors in the conditioning program of the athlete.

- The NATA has a heat illness position statement on their web site (<https://www.nata.org/news-publications/pressroom/statements/position>) with recommendations for prevention: Casa et al. 2015 (<http://natajournals.org/doi/pdf/10.4085/1062-6050-50.9.07>) and Casa & Cisllan, 2009 (<http://natajournals.org/doi/pdf/10.4085/1062-6050-44.3.332>)
- Coaches, athletic trainers, and players should refer to the multiple published best practices by the NATA, American College of Sports Medicine (ACSM), NFHS, and NCAA on preventing and managing heat illness. Emergency action plans should be activated.
- Link to the NFHS Sport Medicine Advisory Committee Position Statements: <https://www.nfhs.org/sports-resource-content/nfhs-sports-medicine-position-statements-and-guidelines/>
- Link to handout from the NATA on Heat Illness: <http://www.nfhs.org/media/1015650/2015-nata-heat-illness-handout.pdf>
- Link to handout from the Kory Stringer Institute on exertional heat stroke prevention: <https://ksi.uconn.edu/wp-content/uploads/sites/1222/2018/01/Preventing-Surviving-EHS-September-2017.pdf>

12. **Head Trauma:** When a player has shown signs or symptoms of head trauma (such as a change in the athlete's behavior, thinking, or physical functioning), the player should receive immediate medical attention from an appropriate medical provider and should not be allowed to return to practice or game that day. The athlete should not be allowed to return to practice or game without an evaluation by an appropriate medical provider.

- All athletes and athletic personnel should follow the state, NFHS, and NCAA policies related to concussion and return to play. See the following CDC resource for a list of states and their concussion policies: <https://www.cdc.gov/headsup/policy/index.html>
- For the most up to date information on concussion management please see the updated Consensus Statement on Concussion in Sport: The 5th International Conference on Concussion in Sport held in Berlin, October 2016 (McCrory et al. 2017 available at <http://bjsm.bmj.com/content/51/11/838>).
- Some cases associated with brain trauma reported that players complained of symptoms or had a previous concussion prior to their deaths. The team physician, athletic trainer, or coach should ensure players understand signs and symptoms of concussion and brain trauma. Players should also be encouraged to inform the team physician, athletic trainer, or coach if they are experiencing any of the signs or symptoms of brain trauma outlined by the CDC.
- **HEADS UP ON CONCUSSION IN SPORTS:**
Information for Parents, Coaches, and School & Sports Professionals. Available at: <http://www.cdc.gov/headsup/highschoolsports/index.html>

The NFHS Sport Medicine Advisory Committee has developed guidelines for concussion management in sports: <http://www.nfhs.org/media/1014737/suggested-guidelines-for-management-of-a-concussion-in-sports-october-2013-2.pdf>

The NCAA has created several rules to help manage concussion injuries. The NCAA has created a set of best practices that are available in the Sports Medicine Handbook which may be found at: <http://www.ncaapublications.com/>

Every NCAA member school is required to have a concussion-management plan that:

- Requires student-athletes to receive information about the signs and symptoms of concussions. They also are required to sign a waiver that says they are responsible for reporting injuries to the medical staff.
- Mandates that institutions provide a process for removing a student-athlete from play/participation if they exhibit signs of a concussion. Student-athletes exhibiting signs of a concussions must be evaluated by a medical staff member with experience in the evaluation and management of concussions before they return to play.

- Prohibits a student-athlete with concussion symptoms from returning to play on the same day of the activity.
- Requires student-athletes diagnosed with a concussion be cleared by a physician before they are permitted to return.

13. **Spinal injuries:** Early recognition, prompt medical evaluation and management of cervical cord and spine injuries is critical for preventing permanent disability and death. Certified athletic trainers are trained to recognize and manage these injuries and whenever possible should be present for all football practices and games. For the most up to date information on management and prevention of these injuries see the following websites:

- National Athletic Trainers Association: <https://www.nata.org/practice-patient-care/health-issues/spine-injury>
- Kory Stringer Institute: <https://ksi.uconn.edu/emergency-conditions/cervical-spine-injury/>
- See also Swartz et al. 2009 for information about cervical spinal injury management and prevention:

<http://www.nata.org/sites/default/files/AcuteMgmtOfCervicalSpineInjuredAthlete.pdf>

14. **Internal Organ Injuries:** Like cervical cord and spinal injuries, early recognition and prompt medical evaluation and treatment of internal organ injuries is critical for ensuring the best possible outcome. Emergency action plans, access to certified athletic trainers, and on-site medical services for competitions constitute best practices for these injuries. A better understanding of the activities and mechanisms associated with these injuries and use of protective gear worn is needed for prevention. Wearing protective gear (e.g., padded belt or shirt) that extends beyond the bottom of the shoulder pads to cover the torso may protect internal organs from direct contact.

CASE SUMMARIES AY2017/18

* Case summaries have been redacted from the publicly available report. Please email nccsir@unc.edu to request case summaries.

REFERENCES

Casa, D., & Csillan, D. (2009). Preseason heat-acclimatization guidelines for secondary school athletics. *Journal of Athletic Training*, 44(3), 332-333. doi: 10.4085/1062-6050-44.3.332

Casa, D. J., J. K. DeMartini, M. F. Bergeron, D. Csillan, E. R. Eichner, R. M. Lopez, M. S. Ferrara, K. C. Miller, F. G. O'Connor, M. N. Sawka and S. W. Yeargin (2015). "National Athletic Trainers' Association position statement: Exertional heat illness." *Journal of Athletic Training* 50(9): 986-1000.

Casa, D., Guskiewicz, K., Anderson, S., Courson, R., Heck, J., Jimenez, C., et al. (2012). National athletic trainers' association position statement: preventing sudden death in sports. *Journal of Athletic Training* 47(1), 96-118.

Drezner, J. A., Courson, R. W., Roberts, W. O., Mosesso, V. N., Link, M. S., & Maron, B. J. (2007). Inter-Association Task Force Recommendations on Emergency Preparedness and Management of Sudden Cardiac Arrest in High School and College Athletic Programs: A Consensus Statement. *Journal of Athletic Training*, 42(1), 143–158.

McCrory, P., W. Meeuwisse, J. Dvořák, M. Aubry, J. Bailes, S. Broglio, et al. (2017). "Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016." *British Journal of Sports Medicine* 51(11): 838-847.

Swartz, E. E., Boden, B. P., Courson, R. W., Decoster, L. C., Horodyski, M., Norkus, S. A., Rehnberg, R. S., Waninger, K. N. (2009). National Athletic Trainers' Association Position Statement: Acute Management of the Cervical Spine–Injured Athlete. *Journal of Athletic Training*, 44(3), 306–331.

Table 1: Number of All catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)

	Collegiate/		High School		All	
	N	%	N	%	N	%
1982-1983	11	18.0%	50	82.0%	61	100.0%
1983-1984	13	21.0%	49	79.0%	62	100.0%
1984-1985	9	17.6%	42	82.4%	51	100.0%
1985-1986	16	27.6%	42	72.4%	58	100.0%
1986-1987	18	25.7%	52	74.3%	70	100.0%
1987-1988	15	17.9%	69	82.1%	84	100.0%
1988-1989	17	23.0%	57	77.0%	74	100.0%
1989-1990	10	13.3%	65	86.7%	75	100.0%
1990-1991	15	24.2%	47	75.8%	62	100.0%
1991-1992	11	23.4%	36	76.6%	47	100.0%
1992-1993	9	15.0%	51	85.0%	60	100.0%
1993-1994	11	16.9%	54	83.1%	65	100.0%
1994-1995	12	23.5%	39	76.5%	51	100.0%
1995-1996	8	15.4%	44	84.6%	52	100.0%
1996-1997	9	13.6%	57	86.4%	66	100.0%
1997-1998	15	20.0%	60	80.0%	75	100.0%
1998-1999	10	12.8%	68	87.2%	78	100.0%
1999-2000	9	12.9%	61	87.1%	70	100.0%
2000-2001	17	24.6%	52	75.4%	69	100.0%
2001-2002	14	17.3%	67	82.7%	81	100.0%
2002-2003	16	25.8%	46	74.2%	62	100.0%
2003-2004	18	24.3%	56	75.7%	74	100.0%
2004-2005	9	13.4%	58	86.6%	67	100.0%
2005-2006	12	21.8%	43	78.2%	55	100.0%
2006-2007	13	17.3%	62	82.7%	75	100.0%
2007-2008	15	18.3%	67	81.7%	82	100.0%
2008-2009	17	15.7%	91	84.3%	108	100.0%
2009-2010	27	27.6%	71	72.4%	98	100.0%
2010-2011	15	20.0%	60	80.0%	75	100.0%
2011-2012	17	21.0%	64	79.0%	81	100.0%
2012-2013	16	32.7%	33	67.3%	49	100.0%
2013-2014	17	21.5%	62	78.5%	79	100.0%
2014-2015	16	19.5%	66	80.5%	82	100.0%
2015-2016	20	19.0%	85	81.0%	105	100.0%
2016-2017	23	29.1%	56	70.9%	79	100.0%
2017-2018	19	23.2%	65	77.4%	84	100.0%
Total	519	20.2%	2047	79.8%	2566	100.0%
Total*	554	20.6%	2132	79.4%	2686	100.0%

*Includes Cheerleading, Drill Team, Rodeo

Table 2: Number of Direct catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)

	Collegiate/		High School		All	
	N	%	N	%	N	%
1982-1983	5	12.5%	35	87.5%	40	100.0%
1983-1984	8	19.0%	34	81.0%	42	100.0%
1984-1985	9	22.5%	31	77.5%	40	100.0%
1985-1986	15	30.6%	34	69.4%	49	100.0%
1986-1987	14	26.9%	38	73.1%	52	100.0%
1987-1988	8	12.3%	57	87.7%	65	100.0%
1988-1989	13	23.2%	43	76.8%	56	100.0%
1989-1990	7	13.5%	45	86.5%	52	100.0%
1990-1991	11	28.2%	28	71.8%	39	100.0%
1991-1992	6	17.6%	28	82.4%	34	100.0%
1992-1993	7	17.9%	32	82.1%	39	100.0%
1993-1994	6	14.0%	37	86.0%	43	100.0%
1994-1995	9	23.7%	29	76.3%	38	100.0%
1995-1996	6	17.6%	28	82.4%	34	100.0%
1996-1997	7	14.0%	43	86.0%	50	100.0%
1997-1998	6	12.0%	44	88.0%	50	100.0%
1998-1999	10	18.2%	45	81.8%	55	100.0%
1999-2000	9	20.9%	34	79.1%	43	100.0%
2000-2001	12	28.6%	30	71.4%	42	100.0%
2001-2002	5	9.6%	47	90.4%	52	100.0%
2002-2003	10	26.3%	28	73.7%	38	100.0%
2003-2004	12	22.2%	42	77.8%	54	100.0%
2004-2005	5	15.6%	27	84.4%	32	100.0%
2005-2006	7	21.9%	25	78.1%	32	100.0%
2006-2007	7	14.9%	40	85.1%	47	100.0%
2007-2008	9	16.7%	45	83.3%	54	100.0%
2008-2009	9	11.5%	69	88.5%	78	100.0%
2009-2010	16	27.1%	43	72.9%	59	100.0%
2010-2011	9	18.4%	40	81.6%	49	100.0%
2011-2012	9	16.4%	46	83.6%	55	100.0%
2012-2013	7	35.0%	13	65.0%	20	100.0%
2013-2014	2	6.1%	31	93.9%	33	100.0%
2014-2015	6	20.7%	23	79.3%	29	100.0%
2015-2016	15	23.8%	48	76.2%	63	100.0%
2016-2017	6	20.7%	23	79.3%	29	100.0%
2017-2018	8	20.5%	31	79.5%	39	100.0%
Total	310	19.1%	1316	80.9%	1626	100.0%
Total*	345	19.9%	1389	80.1%	1734	100.0%

*Includes Cheerleading, Drill Team, Rodeo

Table 3: Number of Indirect catastrophic injuries/illnesses by year: All sports combined, all levels (high school and college)

	Collegiate/		High School		All	
	N	%	N	%	N	%
1982-1983	6	28.6%	15	71.4%	21	100.0%
1983-1984	5	25.0%	15	75.0%	20	100.0%
1984-1985	0	0.0%	11	100.0%	11	100.0%
1985-1986	1	11.1%	8	88.9%	9	100.0%
1986-1987	4	22.2%	14	77.8%	18	100.0%
1987-1988	7	36.8%	12	63.2%	19	100.0%
1988-1989	4	22.2%	14	77.8%	18	100.0%
1989-1990	3	13.0%	20	87.0%	23	100.0%
1990-1991	4	17.4%	19	82.6%	23	100.0%
1991-1992	5	38.5%	8	61.5%	13	100.0%
1992-1993	2	9.5%	19	90.5%	21	100.0%
1993-1994	5	22.7%	17	77.3%	22	100.0%
1994-1995	3	23.1%	10	76.9%	13	100.0%
1995-1996	2	11.1%	16	88.9%	18	100.0%
1996-1997	2	12.5%	14	87.5%	16	100.0%
1997-1998	9	36.0%	16	64.0%	25	100.0%
1998-1999	0	0.0%	23	100.0%	23	100.0%
1999-2000	0	0.0%	27	100.0%	27	100.0%
2000-2001	5	18.5%	22	81.5%	27	100.0%
2001-2002	9	31.0%	20	69.0%	29	100.0%
2002-2003	6	25.0%	18	75.0%	24	100.0%
2003-2004	6	30.0%	14	70.0%	20	100.0%
2004-2005	4	11.4%	31	88.6%	35	100.0%
2005-2006	5	21.7%	18	78.3%	23	100.0%
2006-2007	6	21.4%	22	78.6%	28	100.0%
2007-2008	6	21.4%	22	78.6%	28	100.0%
2008-2009	8	26.7%	22	73.3%	30	100.0%
2009-2010	11	28.2%	28	71.8%	39	100.0%
2010-2011	6	23.1%	20	76.9%	26	100.0%
2011-2012	8	30.8%	18	69.2%	26	100.0%
2012-2013	9	31.0%	20	69.0%	29	100.0%
2013-2014	15	32.6%	31	67.4%	46	100.0%
2014-2015	10	18.9%	43	81.1%	53	100.0%
2015-2016	5	11.9%	37	88.1%	42	100.0%
2016-2017	17	34.0%	33	66.0%	50	100.0%
2017-2018	11	25.6%	34	75.6%	43	100.0%
Total	209	22.3%	731	77.8%	940	100.0%
Total*	209	22.0%	743	78.0%	952	100.0%

*Includes Cheerleading, Drill Team, Rodeo

Table 4a: Number of Direct catastrophic injuries/illnesses by severity by sport: High school all years combined

		Serious		Non-fatal		Fatal		Unknown		All	
		N	%	N	%	N	%	N	%	N	%
Baseball	Male	28	42.4%	21	31.8%	15	22.7%	2	3.0%	66	100.0%
Basketball	Female	3	50.0%	3	50.0%	0	0.0%	0	0.0%	6	100.0%
	Male	9	60.0%	4	26.7%	1	6.7%	1	6.7%	15	100.0%
Cheerleading	Female	42	59.2%	23	32.4%	1	1.4%	5	7.0%	71	100.0%
	Male	1	50.0%	1	50.0%	0	0.0%	0	0.0%	2	100.0%
Cross Country	Female	0	0.0%	0	0.0%	2	100.0%	0	0.0%	2	100.0%
	Male	0	0.0%	2	66.7%	1	33.3%	0	0.0%	3	100.0%
Field Hockey	Female	0	0.0%	1	33.3%	0	0.0%	2	66.7%	3	100.0%
Football	Male	403	41.5%	411	42.3%	137	14.1%	20	2.1%	971	100.0%
Golf	Male	1	100.0%	0	0.0%	0	0.0%	0	0.0%	1	100.0%
Gymnastics	Female	4	36.4%	7	63.6%	0	0.0%	0	0.0%	11	100.0%
	Male	1	25.0%	2	50.0%	1	25.0%	0	0.0%	4	100.0%
Ice Hockey	Female	2	66.7%	1	33.3%	0	0.0%	0	0.0%	3	100.0%
	Male	11	37.9%	14	48.3%	4	13.8%	0	0.0%	29	100.0%
Lacrosse	Female	2	66.7%	0	0.0%	0	0.0%	1	33.3%	3	100.0%
	Male	9	50.0%	6	33.3%	2	11.1%	1	5.6%	18	100.0%
Skiing	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Soccer	Female	5	62.5%	1	12.5%	2	25.0%	0	0.0%	8	100.0%
	Male	7	41.2%	2	11.8%	7	41.2%	1	5.9%	17	100.0%
Softball	Female	6	85.7%	1	14.3%	0	0.0%	0	0.0%	7	100.0%
Swimming	Female	1	16.7%	5	83.3%	0	0.0%	0	0.0%	6	100.0%
	Male	3	30.0%	6	60.0%	1	10.0%	0	0.0%	10	100.0%
Track and Field	Female	7	70.0%	2	20.0%	1	10.0%	0	0.0%	10	100.0%
	Male	14	25.9%	17	31.5%	21	38.9%	2	3.7%	54	100.0%
Volleyball	Male	0	0.0%	1	100.0%	0	0.0%	0	0.0%	1	100.0%
Wrestling	Male	24	35.8%	39	58.2%	3	4.5%	1	1.5%	67	100.0%

Table 4b: Number of Indirect catastrophic injuries/illnesses by severity by sport: High school all years combined

		Serious		Non-fatal		Fatal		Unknown		All	
		N	%	N	%	N	%	N	%	N	%
Baseball	Male	5	19.2%	0	0.0%	21	80.8%	0	0.0%	26	100.0%
Basketball	Female	2	11.1%	0	0.0%	16	88.9%	0	0.0%	18	100.0%
	Male	29	17.8%	0	0.0%	133	82.1%	1	0.6%	163	100.0%
Cheerleading	Female	4	33.3%	0	0.0%	8	66.7%	0	0.0%	12	100.0%
Cross Country	Female	4	28.6%	0	0.0%	10	71.4%	0	0.0%	14	100.0%
	Male	5	19.2%	0	0.0%	21	80.8%	0	0.0%	26	100.0%
	Unknown	0	0.0%	0	0.0%	0	0.0%	1	100.0%	1	100.0%
Field Hockey	Female	0	0.0%	0	0.0%	2	100.0%	0	0.0%	2	100.0%
Football	Male	40	14.1%	2	0.7%	241	84.6%	2	0.7%	285	100.0%
Ice Hockey	Male	3	37.5%	0	0.0%	5	62.5%	0	0.0%	8	100.0%
Lacrosse	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
	Male	1	9.1%	0	0.0%	10	90.9%	0	0.0%	11	100.0%
Other	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Rowing	Male	1	100.0%	0	0.0%	0	0.0%	0	0.0%	1	100.0%
Running/Jogging	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Soccer	Female	2	15.4%	0	0.0%	11	84.6%	0	0.0%	13	100.0%
	Male	8	19.5%	1	2.4%	30	73.2%	2	4.9%	41	100.0%
Softball	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Swimming	Female	2	18.2%	0	0.0%	8	72.7%	1	9.1%	11	100.0%
	Male	1	12.5%	0	0.0%	7	87.5%	0	0.0%	8	100.0%
Tennis	Female	0	0.0%	0	0.0%	2	100.0%	0	0.0%	2	100.0%
	Male	0	0.0%	0	0.0%	4	100.0%	0	0.0%	4	100.0%
Track and Field	Female	0	0.0%	1	14.3%	6	85.7%	0	0.0%	7	100.0%
	Male	6	14.0%	0	0.0%	37	86.0%	0	0.0%	43	100.0%
Volleyball	Female	2	66.7%	0	0.0%	1	33.3%	0	0.0%	3	100.0%
Water Polo	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
	Male	0	0.0%	0	0.0%	3	100.0%	0	0.0%	3	100.0%
Wrestling	Male	5	13.9%	0	0.0%	30	83.3%	1	2.8%	36	100.0%

Table 5a: Number of Direct catastrophic injuries/illnesses by severity by sport: College all years combined

		Serious		Non-fatal		Fatal		Unknown		All	
		N	%	N	%	N	%	N	%	N	%
Baseball	Male	9	45.0%	7	35.0%	3	15.0%	1	5.0%	20	100.0%
Basketball	Male	9	81.8%	1	9.1%	1	9.1%	0	0.0%	11	100.0%
Cheerleading	Female	13	46.4%	13	46.4%	1	3.6%	1	3.6%	28	100.0%
	Male	3	60.0%	2	40.0%	0	0.0%	0	0.0%	5	100.0%
Equestrian	Female	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Field Hockey	Female	2	66.7%	0	0.0%	0	0.0%	1	33.3%	3	100.0%
Football	Male	145	69.0%	48	22.9%	17	8.1%	0	0.0%	210	100.0%
Gymnastics	Female	0	0.0%	2	100.0%	0	0.0%	0	0.0%	2	100.0%
	Male	2	66.7%	1	33.3%	0	0.0%	0	0.0%	3	100.0%
	Unknown	0	0.0%	1	100.0%	0	0.0%	0	0.0%	1	100.0%
Ice Hockey	Female	1	100.0%	0	0.0%	0	0.0%	0	0.0%	1	100.0%
	Male	7	58.3%	5	41.7%	0	0.0%	0	0.0%	12	100.0%
Lacrosse	Female	0	0.0%	2	100.0%	0	0.0%	0	0.0%	2	100.0%
	Male	1	16.7%	1	16.7%	4	66.7%	0	0.0%	6	100.0%
Rodeo	Male	0	0.0%	0	0.0%	2	100.0%	0	0.0%	2	100.0%
Rowing	Male	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Rugby	Male	1	33.3%	2	66.7%	0	0.0%	0	0.0%	3	100.0%
Skiing	Female	0	0.0%	1	50.0%	1	50.0%	0	0.0%	2	100.0%
	Male	0	0.0%	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Soccer	Female	2	50.0%	2	50.0%	0	0.0%	0	0.0%	4	100.0%
	Male	2	66.7%	1	33.3%	0	0.0%	0	0.0%	3	100.0%
Softball	Female	3	75.0%	0	0.0%	0	0.0%	1	25.0%	4	100.0%
Swimming	Male	0	0.0%	1	100.0%	0	0.0%	0	0.0%	1	100.0%
Track and Field	Female	1	50.0%	1	50.0%	0	0.0%	0	0.0%	2	100.0%
	Male	2	14.3%	6	42.9%	6	42.9%	0	0.0%	14	100.0%
Wrestling	Male	1	33.3%	2	66.7%	0	0.0%	0	0.0%	3	100.0%

Table 5b: Number of Indirect catastrophic injuries/illnesses by severity by sport: College all years combined

		Serious		Fatal		Unknown		All	
		N	%	N	%	N	%	N	%
Baseball	Male	2	25.0%	5	62.5%	1	12.5%	8	100.0%
Basketball	Female	3	37.5%	5	62.5%	0	0.0%	8	100.0%
	Male	17	32.1%	36	67.9%	0	0.0%	53	100.0%
Cross Country	Female	0	0.0%	1	100.0%	0	0.0%	1	100.0%
	Male	1	33.3%	2	66.7%	0	0.0%	3	100.0%
Football	Male	13	16.3%	66	82.5%	1	1.3%	80	100.0%
Gymnastics	Female	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Ice Hockey	Male	3	75.0%	1	25.0%	0	0.0%	4	100.0%
Lacrosse	Male	0	0.0%	2	100.0%	0	0.0%	2	100.0%
Rowing	Male	0	0.0%	2	100.0%	0	0.0%	2	100.0%
Skiing	Male	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Soccer	Female	1	25.0%	3	75.0%	0	0.0%	4	100.0%
	Male	2	25.0%	6	75.0%	0	0.0%	8	100.0%
Swimming	Female	1	25.0%	3	75.0%	0	0.0%	4	100.0%
	Male	0	0.0%	8	100.0%	0	0.0%	8	100.0%
Tennis	Female	0	0.0%	1	100.0%	0	0.0%	1	100.0%
	Male	0	0.0%	1	100.0%	0	0.0%	1	100.0%
Track and Field	Male	3	60.0%	2	40.0%	0	0.0%	5	100.0%
Volleyball	Female	2	50.0%	2	50.0%	0	0.0%	4	100.0%
Water Polo	Male	0	0.0%	2	100.0%	0	0.0%	2	100.0%
Wrestling	Male	3	33.3%	6	66.7%	0	0.0%	9	100.0%

Table 6a: Number of Catastrophic injuries/illnesses by Severity by year: High school

	Fatal		Non-fatal		Serious	
	N	Rate per 100,000	N	Rate per 100,000	N	Rate per 100,000
1982-1983	25	0.49	10	0.20	15	0.30
1983-1984	23	0.46	15	0.30	11	0.22
1984-1985	17	0.34	13	0.26	12	0.24
1985-1986	10	0.20	15	0.29	16	0.31
1986-1987	26	0.51	12	0.23	12	0.23
1987-1988	17	0.33	25	0.48	27	0.52
1988-1989	21	0.40	19	0.37	17	0.33
1989-1990	24	0.46	26	0.50	15	0.29
1990-1991	23	0.44	15	0.29	9	0.17
1991-1992	12	0.23	9	0.17	15	0.28
1992-1993	23	0.43	14	0.26	14	0.26
1993-1994	22	0.40	14	0.25	16	0.29
1994-1995	12	0.21	14	0.25	13	0.23
1995-1996	19	0.32	13	0.22	9	0.15
1996-1997	24	0.40	16	0.26	14	0.23
1997-1998	24	0.38	23	0.37	12	0.19
1998-1999	31	0.48	13	0.20	23	0.36
1999-2000	33	0.51	16	0.25	11	0.17
2000-2001	26	0.39	15	0.23	11	0.17
2001-2002	28	0.43	19	0.29	17	0.26
2002-2003	21	0.31	11	0.16	13	0.19
2003-2004	17	0.25	23	0.34	15	0.22
2004-2005	34	0.50	15	0.22	7	0.10
2005-2006	20	0.29	12	0.17	11	0.16
2006-2007	21	0.29	23	0.32	18	0.25
2007-2008	23	0.32	15	0.21	29	0.40
2008-2009	28	0.38	30	0.41	33	0.45
2009-2010	24	0.32	21	0.28	26	0.35
2010-2011	23	0.31	16	0.22	21	0.28
2011-2012	22	0.30	24	0.32	18	0.24
2012-2013	21	0.28	6	0.08	6	0.08
2013-2014	24	0.33	9	0.12	25	0.34
2014-2015	24	0.32	7	0.09	33	0.44
2015-2016	28	0.37	10	0.13	43	0.57
2016-2017	16	0.21	5	0.07	32	0.42
2017-2018	16	0.21	7	0.09	37	0.48

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 6b: Number of Catastrophic injuries/illnesses by Severity by year: College

	Fatal		Non-fatal		Serious	
	N	Rate per 100,000	N	Rate per 100,000	N	Rate per 100,000
1982-1983	7	2.73	3	1.17	1	0.39
1983-1984	5	1.85	2	0.74	6	2.22
1984-1985	1	0.35	3	1.04	5	1.73
1985-1986	4	1.38	4	1.38	8	2.75
1986-1987	5	1.80	2	0.72	11	3.96
1987-1988	7	2.64	1	0.38	7	2.64
1988-1989	4	1.49	4	1.49	9	3.34
1989-1990	3	1.14	2	0.76	4	1.51
1990-1991	5	1.82	6	2.18	4	1.45
1991-1992	5	1.78	1	0.36	5	1.78
1992-1993	3	1.05	0	0.00	6	2.11
1993-1994	7	2.39	0	0.00	4	1.37
1994-1995	3	1.01	3	1.01	6	2.02
1995-1996	2	0.60	4	1.19	2	0.60
1996-1997	3	0.91	5	1.51	1	0.30
1997-1998	9	2.68	1	0.30	5	1.49
1998-1999	2	0.56	3	0.84	5	1.40
1999-2000	2	0.56	2	0.56	5	1.40
2000-2001	7	1.89	4	1.08	6	1.62
2001-2002	10	2.74	1	0.27	3	0.82
2002-2003	6	1.60	3	0.80	7	1.87
2003-2004	9	2.39	4	1.06	5	1.33
2004-2005	4	1.04	1	0.26	4	1.04
2005-2006	5	1.27	4	1.01	2	0.51
2006-2007	6	1.48	1	0.25	6	1.48
2007-2008	5	1.21	1	0.24	9	2.17
2008-2009	5	1.18	1	0.24	11	2.60
2009-2010	13	3.01	2	0.46	12	2.78
2010-2011	4	0.90	7	1.57	3	0.67
2011-2012	9	1.98	0	0.00	8	1.76
2012-2013	8	1.72	3	0.65	4	0.86
2013-2014	6	1.26	0	0.00	11	2.32
2014-2015	4	0.83	1	0.21	11	2.27
2015-2016	3	0.62	3	0.62	13	2.67
2016-2017	5	1.02	1	0.20	17	3.45
2017-2018	5	1.01	1	0.20	13	2.63

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 7a: Rate of direct catastrophic injuries/illnesses by severity by year: High school

	Fatal		Non-fatal		Serious	
	N	Rate per 100,000	N	Rate per 100,000	N	Rate per 100,000
1982-1983	10	0.20	10	0.20	15	0.30
1983-1984	8	0.16	15	0.30	11	0.22
1984-1985	6	0.12	13	0.26	12	0.24
1985-1986	3	0.06	15	0.29	16	0.31
1986-1987	13	0.25	12	0.23	11	0.21
1987-1988	5	0.10	25	0.48	27	0.52
1988-1989	8	0.15	19	0.37	16	0.31
1989-1990	5	0.10	26	0.50	14	0.27
1990-1991	4	0.08	15	0.29	9	0.17
1991-1992	4	0.08	9	0.17	15	0.28
1992-1993	4	0.08	14	0.26	14	0.26
1993-1994	5	0.09	14	0.25	16	0.29
1994-1995	2	0.04	14	0.25	13	0.23
1995-1996	4	0.07	13	0.22	9	0.15
1996-1997	10	0.16	16	0.26	14	0.23
1997-1998	8	0.13	23	0.37	12	0.19
1998-1999	8	0.13	13	0.20	23	0.36
1999-2000	7	0.11	16	0.25	10	0.16
2000-2001	4	0.06	15	0.23	11	0.17
2001-2002	9	0.14	19	0.29	17	0.26
2002-2003	3	0.04	11	0.16	13	0.19
2003-2004	3	0.04	23	0.34	15	0.22
2004-2005	5	0.07	15	0.22	6	0.09
2005-2006	4	0.06	12	0.17	9	0.13
2006-2007	2	0.03	22	0.31	16	0.22
2007-2008	2	0.03	15	0.21	28	0.39
2008-2009	10	0.14	30	0.41	29	0.40
2009-2010	2	0.03	20	0.27	21	0.28
2010-2011	6	0.08	16	0.22	18	0.24
2011-2012	4	0.05	24	0.32	18	0.24
2012-2013	4	0.05	5	0.07	4	0.05
2013-2014	8	0.11	9	0.12	12	0.16
2014-2015	6	0.08	6	0.08	9	0.12
2015-2016	8	0.11	10	0.13	27	0.36
2016-2017	2	0.03	5	0.07	14	0.18
2017-2018	3	0.04	7	0.09	16	0.21

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 7b: Rate of direct catastrophic injuries/illnesses by severity by year: College

	Fatal		Non-fatal		Serious	
	N	Rate per 100,000	N	Rate per 100,000	N	Rate per 100,000
1982-1983	1	0.39	3	1.17	1	0.39
1983-1984	0	0.00	2	0.74	6	2.22
1984-1985	1	0.35	3	1.04	5	1.73
1985-1986	3	1.03	4	1.38	8	2.75
1986-1987	1	0.36	2	0.72	11	3.96
1987-1988	0	0.00	1	0.38	7	2.64
1988-1989	0	0.00	4	1.49	9	3.34
1989-1990	1	0.38	2	0.76	4	1.51
1990-1991	1	0.36	6	2.18	4	1.45
1991-1992	1	0.36	1	0.36	4	1.43
1992-1993	1	0.35	0	0.00	6	2.11
1993-1994	2	0.68	0	0.00	4	1.37
1994-1995	0	0.00	3	1.01	6	2.02
1995-1996	0	0.00	4	1.19	2	0.60
1996-1997	1	0.30	5	1.51	1	0.30
1997-1998	1	0.30	1	0.30	4	1.19
1998-1999	2	0.56	3	0.84	5	1.40
1999-2000	2	0.56	2	0.56	5	1.40
2000-2001	2	0.54	4	1.08	6	1.62
2001-2002	1	0.27	1	0.27	3	0.82
2002-2003	1	0.27	3	0.80	6	1.60
2003-2004	3	0.80	4	1.06	5	1.33
2004-2005	1	0.26	1	0.26	3	0.78
2005-2006	0	0.00	4	1.01	2	0.51
2006-2007	0	0.00	1	0.25	6	1.48
2007-2008	0	0.00	1	0.24	8	1.93
2008-2009	0	0.00	1	0.24	8	1.89
2009-2010	4	0.93	2	0.46	10	2.32
2010-2011	0	0.00	7	1.57	2	0.45
2011-2012	2	0.44	0	0.00	7	1.54
2012-2013	0	0.00	3	0.65	3	0.65
2013-2014	1	0.21	0	0.00	1	0.21
2014-2015	0	0.00	1	0.21	5	1.03
2015-2016	0	0.00	3	0.62	11	2.26
2016-2017	0	0.00	1	0.20	5	1.02
2017-2018	2	0.40	1	0.20	5	1.01

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 8a: Rate of indirect catastrophic injuries/illnesses by severity by year: High School

	Fatal		Non-fatal		Serious	
	N	Rate per 100,000	N	Rate per 100,000	N	Rate per 100,000
1982-1983	15	0.3	0	0	0	0
1983-1984	15	0.3	0	0	0	0
1984-1985	11	0.22	0	0	0	0
1985-1986	7	0.14	0	0	0	0
1986-1987	13	0.25	0	0	1	0.02
1987-1988	12	0.23	0	0	0	0
1988-1989	13	0.25	0	0	1	0.02
1989-1990	19	0.37	0	0	1	0.02
1990-1991	19	0.36	0	0	0	0
1991-1992	8	0.15	0	0	0	0
1992-1993	19	0.36	0	0	0	0
1993-1994	17	0.31	0	0	0	0
1994-1995	10	0.18	0	0	0	0
1995-1996	15	0.25	0	0	0	0
1996-1997	14	0.23	0	0	0	0
1997-1998	16	0.26	0	0	0	0
1998-1999	23	0.36	0	0	0	0
1999-2000	26	0.4	0	0	1	0.02
2000-2001	22	0.33	0	0	0	0
2001-2002	19	0.29	0	0	0	0
2002-2003	18	0.27	0	0	0	0
2003-2004	14	0.21	0	0	0	0
2004-2005	29	0.42	0	0	1	0.01
2005-2006	16	0.23	0	0	2	0.03
2006-2007	19	0.27	1	0.01	2	0.03
2007-2008	21	0.29	0	0	1	0.01
2008-2009	18	0.25	0	0	4	0.05
2009-2010	22	0.3	1	0.01	5	0.07
2010-2011	17	0.23	0	0	3	0.04
2011-2012	18	0.24	0	0	0	0
2012-2013	17	0.23	1	0.01	2	0.03
2013-2014	16	0.22	0	0	13	0.18
2014-2015	18	0.24	1	0.01	24	0.32
2015-2016	20	0.26	0	0	16	0.21
2016-2017	14	0.18	0	0	18	0.23
2017-2018	13	0.17	0	0	21	0.27

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 8b: Rate of indirect catastrophic injuries/illnesses by severity by year: College

	N	Fatal Rate per 100,000	N	Serious Rate per 100,000
1982-1983	6	2.34	0	0
1983-1984	5	1.85	0	0
1985-1986	1	0.34	0	0
1986-1987	4	1.44	0	0
1987-1988	7	2.64	0	0
1988-1989	4	1.49	0	0
1989-1990	2	0.76	0	0
1990-1991	4	1.45	0	0
1991-1992	4	1.43	1	0.36
1992-1993	2	0.7	0	0
1993-1994	5	1.71	0	0
1994-1995	3	1.01	0	0
1995-1996	2	0.6	0	0
1996-1997	2	0.6	0	0
1997-1998	8	2.38	1	0.3
2000-2001	5	1.35	0	0
2001-2002	9	2.47	0	0
2002-2003	5	1.34	1	0.27
2003-2004	6	1.59	0	0
2004-2005	3	0.78	1	0.26
2005-2006	5	1.27	0	0
2006-2007	6	1.48	0	0
2007-2008	5	1.21	1	0.24
2008-2009	5	1.18	3	0.71
2009-2010	9	2.08	2	0.46
2010-2011	4	0.9	1	0.22
2011-2012	7	1.54	1	0.22
2012-2013	8	1.72	1	0.22
2013-2014	5	1.05	10	2.11
2014-2015	4	0.83	6	1.24
2015-2016	3	0.62	2	0.41
2016-2017	5	1.02	12	2.44
2017-2018	3	0.61	8	1.62

Note: Rates with number of incidents less than 5 should be interpreted with caution.

**Table 9a: Rate of Direct catastrophic injuries/illnesses by level and severity by sport:
High school**

		N	Fatal Rate per 100,000	N	Non-fatal Rate per 100,000	N	Serious Rate per 100,000
Baseball	Male	15	0.09	21	0.13	28	0.17
Basketball	Female	0	0	3	0.02	3	0.02
	Male	1	0.01	4	0.02	9	0.05
Cheerleading	Female	1	0.04	23	0.98	42	1.8
	Male	0	0	1	1.71	1	1.71
Cross Country	Female	2	0.04	0	0	0	0
	Male	1	0.02	2	0.03	0	0
Field Hockey	Female	0	0	1	0.05	0	0
Football	Male	137	0.38	411	1.14	403	1.12
Golf	Male	0	0	0	0	1	0.02
Gymnastics	Female	0	0	7	0.85	4	0.48
	Male	1	0.84	2	1.68	1	0.84
Ice Hockey	Female	0	0	1	0.61	2	1.22
	Male	4	0.37	14	1.3	11	1.02
Lacrosse	Female	0	0	0	0	2	0.14
	Male	2	0.11	6	0.32	9	0.48
Skiing	Female	1	0.33	0	0	0	0
Soccer	Female	2	0.02	1	0.01	5	0.06
	Male	7	0.06	2	0.02	7	0.06
Softball	Female	0	0	1	0.01	6	0.05
Swimming	Female	0	0	5	0.11	1	0.02
	Male	1	0.03	6	0.16	3	0.08
Track and Field	Female	1	0.01	2	0.01	7	0.04
	Male	21	0.11	17	0.09	14	0.07
Volleyball	Male	0	0	1	0.08	0	0
Wrestling	Male	3	0.03	39	0.44	24	0.27

Note: Rates with number of incidents less than 5 should be interpreted with caution.

**Table 9b: Rate of Direct catastrophic injuries/illnesses by level and severity by sport:
College**

		N	Fatal Rate per 100,000	N	Non-fatal Rate per 100,000	N	Serious Rate per 100,000
Baseball	Male	3	0.32	7	0.74	9	0.95
Basketball	Male	1	0.18	1	0.18	9	1.62
Cheerleading	Female	1	0	13	0	13	0
	Male	0	0	2	0	3	0
Equestrian	Female	1	3.88	0	0	0	0
Field Hockey	Female	0	0	0	0	2	1.03
Football	Male	17	0.81	48	2.28	145	6.89
Gymnastics	Female	0	0	2	3.72	0	0
	Male	0	0	1	5.37	2	10.74
Ice Hockey	Female	0	0	0	0	1	2.54
	Male	0	0	5	3.56	7	4.98
Lacrosse	Female	0	0	2	0.98	0	0
	Male	4	1.46	1	0.36	1	0.36
Rowing	Male	1	1.28	0	0	0	0
Skiing	Female	1	5.59	1	5.59	0	0
	Male	1	4.62	0	0	0	0
Soccer	Female	0	0	2	0.34	2	0.34
	Male	0	0	1	0.15	2	0.3
Softball	Female	0	0	0	0	3	0.58
Swimming	Male	0	0	1	0.34	0	0
Track and Field	Female	0	0	1	0.08	1	0.08
	Male	6	0.41	6	0.41	2	0.14
Wrestling	Male	0	0	2	0.81	1	0.41

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 10a: Indirect catastrophic events by level and severity: High school

		N	Fatal Rate per 100,000	N	Non-fatal Rate per 100,000	N	Serious Rate per 100,000
Baseball	Male	21	0.13	0	0	5	0.03
Basketball	Female	16	0.1	0	0	2	0.01
	Male	133	0.69	0	0	29	0.15
Cheerleading	Female	8	0.34	0	0	4	0.17
Cross Country	Female	10	0.18	0	0	4	0.07
	Male	21	0.32	0	0	5	0.08
Field Hockey	Female	2	0.1	0	0	0	0
Football	Male	241	0.67	2	0.01	40	0.11
Ice Hockey	Male	5	0.46	0	0	3	0.28
Lacrosse	Female	1	0.07	0	0	0	0
	Male	10	0.53	0	0	1	0.05
Rowing	Male	0	0	0	0	1	1.79
Soccer	Female	11	0.12	0	0	2	0.02
	Male	30	0.26	1	0.01	8	0.07
Softball	Female	1	0.01	0	0	0	0
Swimming	Female	8	0.17	0	0	2	0.04
	Male	7	0.19	0	0	1	0.03
Tennis	Female	2	0.04	0	0	0	0
	Male	4	0.08	0	0	0	0
Track and Field	Female	6	0.04	1	0.01	0	0
	Male	37	0.19	0	0	6	0.03
Volleyball	Female	1	0.01	0	0	2	0.02
Water Polo	Female	1	0.26	0	0	0	0
	Male	3	0.57	0	0	0	0
Wrestling	Male	30	0.34	0	0	5	0.06

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 10b: Indirect catastrophic events by level and severity: College

		N	Fatal Rate per 100,000	N	Serious Rate per 100,000
Baseball	Male	5	0.53	2	0.21
Basketball	Female	5	1.02	3	0.61
	Male	36	6.46	17	3.05
Cross Country	Female	1	0.25	0	0
	Male	2	0.49	1	0.24
Football	Male	66	3.14	13	0.62
Gymnastics	Female	1	1.86	0	0
Ice Hockey	Male	1	0.71	3	2.13
Lacrosse	Male	2	0.73	0	0
Rowing	Male	2	2.57	0	0
Skiing	Male	1	4.62	0	0
Soccer	Female	3	0.51	1	0.17
	Male	6	0.9	2	0.3
Swimming	Female	3	0.84	1	0.28
	Male	8	2.7	0	0
Tennis	Female	1	0.34	0	0
	Male	1	0.36	0	0
Track and Field	Male	2	0.14	3	0.21
Volleyball	Female	2	0.43	2	0.43
Water Polo	Male	2	5.63	0	0
Wrestling	Male	6	2.44	3	1.22

Note: Rates with number of incidents less than 5 should be interpreted with caution.

Table 11: Characteristics of all sport-related catastrophic events during AY 2017-2018

	Direct		Indirect		All	
	N	%	N	%	N	%
Total	39	100.0%	46	100.0%	85	100.0%
Sport Level						
Collegiate/University	8	20.5%	11	23.9%	19	22.4%
High School Sponsored	31	79.5%	35	76.1%	66	77.6%
Severity						
Serious	21	53.8%	30	65.2%	51	60.0%
Non-fatal	8	20.5%	0	0.0%	8	9.4%
Fatal	5	12.8%	16	34.8%	21	24.7%
Unknown	5	12.8%	0	0.0%	5	5.9%
Sex						
Female	3	7.7%	3	6.5%	6	7.1%
Male	36	92.3%	43	93.5%	79	92.9%
Month						
Jul-Aug	5	12.8%	10	21.7%	15	17.6%
Sep-Oct	28	71.8%	13	28.3%	41	48.2%
Nov-Dec	0	0.0%	7	15.2%	7	8.2%
Jan-Feb	2	5.1%	6	13.0%	8	9.4%
Mar-Apr	1	2.6%	4	8.7%	5	5.9%
May-Jun	3	7.7%	6	13.0%	9	10.6%
Sport						
Baseball	2	5.1%	2	4.3%	4	4.7%
Basketball	0	0.0%	13	28.3%	13	15.3%
Cheerleading	0	0.0%	1	2.2%	1	1.2%
Cross Country	1	2.6%	2	4.3%	3	3.5%
Football	31	79.5%	15	32.6%	46	54.1%
Ice Hockey	1	2.6%	0	0.0%	1	1.2%
Lacrosse	1	2.6%	2	4.3%	3	3.5%
Soccer	2	5.1%	6	13.0%	8	9.4%
Swimming	0	0.0%	2	4.3%	2	2.4%
Track and Field	0	0.0%	2	4.3%	2	2.4%
Wrestling	1	2.6%	1	2.2%	2	2.4%
Sponsored activity						
Official school or team related ATHLETIC activity (e.g. official practice, team strength/fitness training or competition)	39	100.0%	37	80.4%	76	89.4%
Personal athletic activity (e.g. individual strength/fitness or practice, non-team related)	0	0.0%	9	19.6%	9	10.6%
Location						
Athlete's Home	0	0.0%	2	4.3%	2	2.4%
Competitive Venue (e.g. arena, stadium, track, field)	33	84.6%	22	47.8%	55	64.7%

	Direct		Indirect		All	
	N	%	N	%	N	%
Other Private Property	0	0.0%	1	2.2%	1	1.2%
Public Park or Street	1	2.6%	1	2.2%	2	2.4%
School Athletic Facility (e.g. weight room, practice field)	5	12.8%	19	41.3%	24	28.2%
School Campus (not at an athletic facility)	0	0.0%	1	2.2%	1	1.2%
Event Type						
Competition/Game	27	69.2%	16	34.8%	43	50.6%
Conditioning Session	1	2.6%	3	6.5%	4	4.7%
Other	0	0.0%	1	2.2%	1	1.2%
Practice	10	25.6%	15	32.6%	25	29.4%
Scrimmage	1	2.6%	3	6.5%	4	4.7%
Strength/Weight Session	0	0.0%	2	4.3%	2	2.4%
Unaffiliated Recreational Activity	0	0.0%	6	13.0%	6	7.1%
Player action						
5,000 m/3mi	0	0.0%	2	4.3%	2	2.4%
Being blocked	2	5.1%	0	0.0%	2	2.4%
Being tackled	5	12.8%	0	0.0%	5	5.9%
Blocking	2	5.1%	1	2.2%	3	3.5%
Conditioning (land)	1	2.6%	10	21.7%	11	12.9%
Conditioning (water)	0	0.0%	1	2.2%	1	1.2%
Defending	1	2.6%	0	0.0%	1	1.2%
Fielding	2	5.1%	0	0.0%	2	2.4%
Fitness - Other	0	0.0%	1	2.2%	1	1.2%
General play	2	5.1%	11	23.9%	13	15.3%
Heading ball	1	2.6%	0	0.0%	1	1.2%
Other	0	0.0%	3	6.5%	3	3.5%
Penalty shot	0	0.0%	1	2.2%	1	1.2%
Receiving pass	1	2.6%	0	0.0%	1	1.2%
Running	1	2.6%	0	0.0%	1	1.2%
Running (middle/long distance)	1	2.6%	0	0.0%	1	1.2%
Running (sprints)	0	0.0%	1	2.2%	1	1.2%
Tackling	9	23.1%	0	0.0%	9	10.6%
Takedown	1	2.6%	0	0.0%	1	1.2%
Throwing	0	0.0%	1	2.2%	1	1.2%
Throwing - discus	0	0.0%	1	2.2%	1	1.2%
Unknown	10	25.6%	12	26.1%	22	25.9%
Weights	0	0.0%	1	2.2%	1	1.2%
Basic Mechanism						
Contact with Another Player	22	56.4%	0	0.0%	22	25.9%
Contact with Apparatus or Object	5	12.8%	0	0.0%	5	5.9%
Contact with Ground/Surface	4	10.3%	0	0.0%	4	4.7%
Environmental (e.g., lightning strike)	0	0.0%	6	13.0%	6	7.1%
Infection or Illness	0	0.0%	37	80.4%	37	43.5%

	Direct		Indirect		All	
	N	%	N	%	N	%
No Direct or Indirect Contact	0	0.0%	2	4.3%	2	2.4%
Other	1	2.6%	1	2.2%	2	2.4%
Unknown	7	17.9%	0	0.0%	7	8.2%
Major Injury Category						
Head Injury	13	33.3%	0	0.0%	13	15.3%
Heat-related injury	0	0.0%	8	17.4%	8	9.4%
Hit in the Chest	2	5.1%	0	0.0%	2	2.4%
Other	0	0.0%	4	8.7%	4	4.7%
Other Traumatic Injury	6	15.4%	0	0.0%	6	7.1%
Spinal Cord Injury	18	46.2%	0	0.0%	18	21.2%
Sudden Cardiac Arrest	0	0.0%	33	71.7%	33	38.8%
Unknown at this time	0	0.0%	1	2.2%	1	1.2%
Detailed Injury Category						
Cardiac/Sudden Cardiac Arrest (not Commotio Cordis)	0	0.0%	33	71.7%	33	38.8%
Commotio Cordis (external blunt chest wall impact resulting in Cardiac Arrest)	2	5.1%	0	0.0%	2	2.4%
Heat-Related Injury (e.g. Heatstroke)	0	0.0%	7	15.2%	7	8.2%
Other	0	0.0%	5	10.9%	5	5.9%
Other Traumatic Injury (e.g. Ruptured Spleen)	7	17.9%	0	0.0%	7	8.2%
Rhabdomyolysis	0	0.0%	1	2.2%	1	1.2%
Spinal Cord Injury with a Fracture	6	15.4%	0	0.0%	6	7.1%
Spinal Cord Injury without Spine Fracture	7	17.9%	0	0.0%	7	8.2%
Spine Fracture	5	12.8%	0	0.0%	5	5.9%
Traumatic Brain Injury (e.g. subdural hematoma)	12	30.8%	0	0.0%	12	14.1%
Injury Outcome						
Fatality/Sudden Death	5	12.8%	16	34.8%	21	24.7%
Non-trauma Survivor (e.g. sudden cardiac arrest, heat stroke, exertional sickling)	0	0.0%	30	65.2%	30	35.3%
Trauma-related Non-Fatality - Disability unknown/uncertain	5	12.8%	0	0.0%	5	5.9%
Trauma-related Non-Fatality with Permanent Disability	8	20.5%	0	0.0%	8	9.4%
Trauma-related Non-Fatality with Temporary Disability (full recovery expected or confirmed)	21	53.8%	0	0.0%	21	24.7%

Table 12. Participation numbers, 1982/83 to 2017/18

	High School ¹		College ²	
	Female	Male	Female	Male
Baseball	34,773	16,144,184	--	950,070
Basketball	15,372,149	19,235,256	491,881	556,893
Cheerleading³	2,338,643	58,423	--	--
Cross Country	5,633,433	6,659,059	406,777	411,912
Equestrian⁴	23,176	3,856	25,790	1,066
Field Hockey	2,046,821	5,220	194,935	--
Football	34,033	36,127,067	--	2,104,403
Golf	1,912,143	5,164,308	110,089	276,796
Gymnastics	825,004	119,240	53,833	18,625
Ice Hockey	163,692	1,080,963	39,439	140,620
Lacrosse	1,393,614	1,886,197	204,678	274,102
Rowing⁴	63,741	55,913	172,888	77,824
Skiing	301,388	360,539	17,875	21,667
Soccer	8,913,763	11,376,702	586,314	667,259
Softball	11,926,428	54,805	514,901	--
Swimming/Diving	4,572,514	3,653,574	357,602	296,230
Tennis	5,609,619	5,210,170	293,098	279,919
Track and Field⁵	16,174,896	19,834,884	1,259,684	1,451,603
Volleyball⁶	13,092,111	1,242,277	462,984	42,520
Water Polo	382,639	525,064	23,792	35,535
Wrestling	148,965	8,849,306	--	245,578

¹NFHS available online: <https://www.nfhs.org/ParticipationStatistics/ParticipationStatistics/>

²NCAA accessed online: https://ncaaorg.s3.amazonaws.com/research/sportpart/Oct2018RES_2017-18SportsSponsorshipParticipationRatesReport.pdf

³Cheerleading is not an official sport for NCAA collegiate athletes.

⁴Equestrian (male and female) and rowing (males) are non-championship NCAA collegiate sports.

⁵Includes both indoor and outdoor track and field.

⁶Includes sand volleyball.


Note: Not all high schools and colleges are members of the NFHS and NCAA. Complete data are not available for the non-member schools. Therefore, these participation numbers underestimate the total number of high school and collegiate participants in the United States.

Figure 1: Rates of fatal catastrophic direct and indirect injuries/illnesses by sport-gender among high school participants, 1982/83-2017/18




Note: Rates with number of incidents less than 5 should be interpreted with caution.

Figure 2: Rates of all catastrophic direct and indirect injuries/illnesses by sport-gender among high school participants, 1982/83-2017/18




Note: Rates with number of incidents less than 5 should be interpreted with caution.

Figure 3: Rates of fatal catastrophic direct and indirect injuries/illnesses by sport-gender among collegiate participants, 1982/83-2017/18



Note: Rates with number of incidents less than 5 should be interpreted with caution.

Figure 4: Rates of all catastrophic direct and indirect injuries/illnesses by sport-gender among collegiate participants, 1982/83-2017/18



Note: Rates with number of incidents less than 5 should be interpreted with caution.